

Helical glasses near ferromagnetic quantum criticality

S. J. Thomson,¹ F. Krüger,¹ and A. G. Green²

¹*SUPA, School of Physics and Astronomy, University of St. Andrews,
North Haugh, St. Andrews, KY16 9SS, United Kingdom*

²*London Centre for Nanotechnology, University College London,
Gordon St., London, WC1H 0AH, United Kingdom*

(Dated: March 20, 2013)

We study the effects of quenched charge disorder on the phase reconstruction near itinerant ferromagnetic quantum critical points in three spatial dimensions. Combining a replica disorder average with a fermionic version of the quantum order-by-disorder mechanism, we show that weak disorder destabilizes the ferromagnetic state and enhances the susceptibility towards incommensurate, spiral magnetic ordering. The Goldstone modes of the spiral phase are governed by a $3d$ -XY model. The induced disorder in the pitch of the spiral generates a random anisotropy for the Goldstone modes, inducing vortex lines in the phase of the helical order and rendering the magnetic correlations short ranged with a strongly anisotropic correlation length.

PACS numbers: 64.70.Tg, 75.30.Kz, 75.10.Lp, 75.10.Nr

Quantum criticality remains one of the great enigmas of modern condensed matter physics. Fluctuations around quantum critical points are now known to be responsible for many unexpected phenomena, such as the discontinuous quantum phase transitions seen in many itinerant ferromagnets (FM). Examples include MnSi [1, 2], $\text{Sr}_{1-x}\text{Ca}_x\text{RuO}_3$ [2], CoO_2 [3], UGe_2 [4], and URhGe [5]. Such generic fluctuation-induced first-order behavior is a consequence of the coupling between the magnetic order-parameter- and soft electronic particle-hole fluctuations [6, 7].

Fluctuation-induced first-order behavior can be indicative of potential instabilities towards incommensurate ordering [7]. Recently, this possibility has been explored within a fermionic version of the quantum order-by-disorder mechanism [8–10]. The key idea is that certain deformations of the Fermi surface associated with the onset of competing order enhance the phase space available for low-energy particle-hole fluctuations. The coupling of these additional soft modes to the order parameter fluctuations leads to a self-consistent stabilization of incommensurate, spiral magnetic order, pre-empting the first-order transition to the homogenous FM.

While disorder can smear continuous quantum phase transitions due to quantum Griffiths effects [11, 12], the finite temperature tricritical point, below which the magnetic transition turns first order and spiral order is predicted, survives up to a critical disorder strength [6]. In this Letter, we argue that the interplay between disorder and quantum fluctuations leads to a helical glass state consistent with the strongly inhomogeneous short-ranged ordered state observed recently near the avoided FM quantum critical point of CeFePO [13].

We consider electrons in three dimensions at chemical potential μ , interacting via a local repulsion g , and subject to quenched charge disorder. The disorder potential $v_c(\mathbf{r})$ is Gaussian distributed with zero mean, variance σ_c^2

and uncorrelated between different positions. The Hamiltonian expressed in terms of the fermionic field operators $\psi(\mathbf{r}) = (\psi_\uparrow(\mathbf{r}), \psi_\downarrow(\mathbf{r}))$, is given by

$$\mathcal{H} = \int d^3\mathbf{r} \left\{ \psi^\dagger [\nabla^2 - \mu + v_c(\mathbf{r})] \psi + g \psi_\uparrow^\dagger \psi_\uparrow \psi_\downarrow^\dagger \psi_\downarrow \right\}. \quad (1)$$

Our discussion of this model is divided into four parts. First, we summarize the phase diagram of the clean itinerant FM. A fermionic quantum order-by-disorder treatment of fluctuations shows that FM order gives way to fluctuation-driven spiral order in the vicinity of the quantum critical point. Next, we treat the effects of weak disorder using the replica trick alongside the fermionic order-by-disorder approach. We find that although FM order is destabilized by disorder, the region of the phase diagram occupied by spiral order is enhanced. Next, we turn to the effects of disorder within the spiral phase. Taking account of lattice-induced anisotropy in the orientation of the spiral wave-vector and the combination of this with disorder, we show that the Goldstone modes are described by a $3d$ -XY model with random anisotropy. This form of disorder has a dramatic effect, driving the formation of vortex lines in the phase of the spiral and resulting in glassy behavior with short range magnetic correlations. Finally, we discuss our results in the context of recent experiments [13] and blue phases and lines of skyrmion defects predicted by other authors [14, 15].

Clean System: The free energy of the clean system described by (1) for $v_c(\mathbf{r}) = 0$ can, in mean field theory, be conveniently expressed in a Ginzburg-Landau expansion about the paramagnetic state: $F_{\text{mf}}(m) \approx am^2 + bm^4 + cm^6$, with coefficients that are known functions of temperature and interaction strength. Restricting for a moment to a spatially uniform FM, the fluctuation correction to the free energy takes the form [6–10] $F_{\text{fl}}(m) = \frac{1}{2} \lambda \tilde{g}^2 m^4 \ln(\kappa m^2 + \tilde{T}^2)$, where λ and κ are constants [16]. Here and in the following we use rescaled,




FIG. 1: (color online) Fluctuations near the FM quantum critical point stabilize an inhomogeneous, spiral magnetic state. While disorder ($\tilde{\sigma}_c^2 = 0.4$) destabilizes the FM, it enhances the region of spiral order.

dimensionless units, $\tilde{T} = T/\mu$ and $\tilde{g} = g\rho_F$ (ρ_F is the density of states at the Fermi level). This correction to the free energy shows a $\ln T$ -divergence of the m^4 coefficient, ultimately driving the transition first order.

In fact, it has become apparent from subsequent analysis [8–10] that the first order FM transition is pre-empted by a transition into a spiral phase with order parameter $\mathbf{m}(\mathbf{r}) = m[\mathbf{n}_x \cos(\mathbf{q} \cdot \mathbf{r}) + \mathbf{n}_y \sin(\mathbf{q} \cdot \mathbf{r})]$ with $\mathbf{q} = q\mathbf{n}_z$. For a spiral of pitch q (measured in units of the Fermi momentum) we may use the fact that the free energy is a functional of the mean-field electron dispersion - $\epsilon_\nu(\mathbf{k}) = k^2 + \nu\sqrt{(k_z q)^2 + m^2}$ - together with the fact that the important integrals are peaked near the Fermi surface at low temperatures, to write the free energy as an angular average over $\hat{k}_z = k_z/k$, $F(m, q) = \langle F(\sqrt{k_z^2 q^2 + m^2}) \rangle$.

This results in relationships between powers of m and q in the Ginzburg-Landau expansion, e.g. the $m^2 q^2$ coefficient is proportional to the logarithmically divergent m^4 coefficient, explaining why the spiral is stabilized below some temperature. We expand in powers of q , $F(m, q) = f_0(m^2) + f_2(m^2)q^2 + \frac{1}{2}f_4(m^2)q^4$, but keep the full functional forms of the functions $f_{2n}(m^2)$. The phase diagram shown in Fig. 1 is obtained by minimizing the resulting free energy $F(m, q)$.

Replica Treatment of Weak Disorder: The effect of disorder on the magnetic phase diagram, can be addressed using a combination of quantum order-by-disorder with the replica trick. We first recapitulate the steps involved in deriving the fluctuation corrections and incorporate disorder at the appropriate stage. This proceeds in four steps: (i) decouple the interaction by a Hubbard-Stratonovich transformation in spin and charge channels.

(ii) separate zero- and finite-frequency spin and charge fluctuations, allowing the possibility of a static magnetic spiral background. (iii) integrate out the electrons treating disorder perturbatively to leading order. The latter introduces new terms in the free energy not found for the clean system. (iv) integrate out spin and charge fluctuations to quadratic order to obtain an effective Ginzburg-Landau description of the magnetic phase diagram.

(i) The Hubbard-Stratonovich transformation is made by introducing classical fields $\rho(\mathbf{r}, \tau)$ and $\phi(\mathbf{r}, \tau)$, representing charge and spin, respectively. In order to compute the disorder averaged free energy we employ the replica trick, $\bar{F} = -T \lim_{n \rightarrow 0} (\bar{\mathcal{Z}}^n - 1)/n$. The effective replica action S_n , defined by $\bar{\mathcal{Z}}^n = \int \mathcal{D}[\bar{\psi}^\alpha, \psi^\alpha, \rho^\alpha, \phi^\alpha] \exp(-S_n)$, consists of n copies of the clean system and a disorder vertex which is non-diagonal in the replica index and in imaginary time,

$$S_n = \sum_{\alpha=1}^n \int d\tau \int d^3\mathbf{r} \left\{ g \left[(\phi^\alpha)^2 - (\rho^\alpha)^2 \right] + \bar{\psi}^\alpha \left[\partial_\tau - \nabla^2 - \mu + g(\rho^\alpha - \phi^\alpha \cdot \boldsymbol{\sigma}) \right] \psi^\alpha \right\} - \frac{\sigma_c^2}{2} \sum_{\alpha, \alpha'} \sum_{\nu, \nu' = \pm 1} \int_{\tau, \tau'} \int_{\mathbf{r}} \bar{\psi}_\nu^\alpha(\tau) \psi_\nu^\alpha(\tau) \bar{\psi}_{\nu'}^{\alpha'}(\tau') \psi_{\nu'}^{\alpha'}(\tau'). \quad (2)$$

We have introduced the vector $\boldsymbol{\sigma} = (\sigma^x, \sigma^y, \sigma^z)^T$ of Pauli matrices to conveniently express the electron spin.

(ii) We decompose the fields ρ and ϕ into zero- and finite-frequency components, the former corresponding to static order in the system, $\rho^\alpha(\mathbf{r}, \tau) = \rho_0(\mathbf{r}) + \tilde{\rho}^\alpha(\mathbf{r}, \tau)$ and $\phi^\alpha(\mathbf{r}, \tau) = \mathbf{m}(\mathbf{r}) + \tilde{\phi}^\alpha(\mathbf{r}, \tau)$. In the following, we do not consider the possibility of charge order ($\rho_0 = 0$) and specify $\mathbf{m}(\mathbf{r})$ to be the spiral order parameter.

Including $\mathbf{m}(\mathbf{r})$ in the free-fermion action allows a self-consistent free energy expansion in the presences of spiral magnetic order. Transforming the fermions to the rotating spiral frame,

$$\varphi_\nu^\alpha(\mathbf{k}, \omega_n) = \sum_{\nu'} \left(e^{-i\frac{\theta(\mathbf{k})}{2}\sigma_y} \right)_{\nu, \nu'} \psi_{\nu'}^\alpha(\mathbf{k} + \nu' \frac{\mathbf{q}}{2}, \omega_n), \quad (3)$$

with $\cot \theta(\mathbf{k}) = k_z q/m$ and ω_n denoting fermionic Matsubara frequencies, we obtain a diagonal free-fermion action $S_0[\bar{\varphi}, \varphi, \mathbf{m}]$ with corresponding Green's function $G_\nu(\mathbf{k}, \omega_n) = (-i\omega_n + \epsilon_\nu(\mathbf{k}) - \mu)^{-1}$.

(iii) Next, we integrate over the fermion fields $\bar{\varphi}, \varphi$, treating the disorder vertex $S_{\text{dis}}[\bar{\varphi}, \varphi]$ perturbatively to leading order. The resulting mean-field action $S_{\text{mf}}[\mathbf{m}]$, which is independent of the fluctuation fields, is given by

$$S_{\text{mf}} = n \frac{gm^2}{T} - n \sum_{\nu \omega_n} \int_{\mathbf{k}} \ln G_\nu^{-1}(\mathbf{k}, \omega_n) + \langle S_{\text{dis}}[\bar{\varphi}, \varphi] \rangle_0. \quad (4)$$

The average $\langle \dots \rangle_0$ is taken with respect to the free-fermion action $S_0[\bar{\varphi}, \varphi, \mathbf{m}]$ in the presence of spiral order. The fluctuation corrections, up to quadratic order in the

fluctuation fields, are given by

$$\begin{aligned} \tilde{S}_{\text{fl}}[\mathbf{m}, \tilde{\rho}, \tilde{\phi}] &= \tilde{S}_0[\tilde{\rho}, \tilde{\phi}] - \frac{1}{2} \left\langle \tilde{S}^2[\tilde{\varphi}, \varphi, \tilde{\rho}, \tilde{\phi}] \right\rangle_0 \\ &+ \frac{1}{2} \left\langle \tilde{S}^2[\tilde{\varphi}, \varphi, \tilde{\rho}, \tilde{\phi}] S_{\text{dis}}[\tilde{\varphi}, \varphi] \right\rangle_0^{(c)} \end{aligned} \quad (5)$$

where $\langle \tilde{S}^2 S_{\text{dis}} \rangle^{(c)} = \langle \tilde{S}^2 S_{\text{dis}} \rangle - \langle \tilde{S}^2 \rangle \langle S_{\text{dis}} \rangle$ only includes connected diagrams. Here $\tilde{S}_0[\tilde{\rho}, \tilde{\phi}]$ is the contribution arising from the first line of Eq. (2) while $\tilde{S}[\tilde{\varphi}, \varphi, \tilde{\rho}, \tilde{\phi}]$ denotes the linear coupling of the fluctuations to the fermion fields originating from the second line of Eq. (2).

(iv) To obtain the fluctuation corrections to the disorder averaged free energy, we perform the Gaussian integrals over the fluctuation fields, re-exponentiate, and finally take the replica limit $n \rightarrow 0$.

Including disorder modifies both the mean field and fluctuation corrections to the free energy. The proportionality between expansion coefficients found in the clean system is broken in the presence of disorder since q enters not only the electron dispersion, but also through the transformation (3) of the disorder vertex. The mean-field free energy contribution due to disorder is given by

$$F_{\text{mf}}^{\text{dis}} = \frac{\tilde{\sigma}_c^2}{2} \sum_{\nu, \nu'} \int_{\mathbf{k}, \mathbf{k}'} h_{\nu, \nu'}(\mathbf{k}, \mathbf{k}') \chi_{\nu, \nu'}(\mathbf{k}, \mathbf{k}'), \quad (6)$$

where $\chi_{\nu, \nu'}(\mathbf{k}, \mathbf{k}') = \tilde{T} \sum_{\omega_n} G_{\nu}(\mathbf{k}, \omega_n) G_{\nu'}(\mathbf{k}', \omega_n) = [n_F(\epsilon_{\nu}(\mathbf{k})) - n_F(\epsilon_{\nu'}(\mathbf{k}'))]/[\epsilon_{\nu}(\mathbf{k}) - \epsilon_{\nu'}(\mathbf{k}')] with $n_F(\epsilon) = 1/[\exp(\frac{\epsilon-1}{\tilde{T}}) + 1]$ the Fermi-function and $h_{\nu, -\nu} = 1 - h_{\nu, \nu} \approx \frac{1}{4}(k_z - k'_z)^2 q^2/m^2$. The disorder strength in dimensionless units is given by $\tilde{\sigma}_c^2 = \sigma_c^2 \rho_F/\mu$.$

We have expanded $h_{\nu, \nu'}$ up to quadratic order in q , since we will only compute disorder corrections to the m^2 , m^4 , and $q^2 m^2$ coefficients. Spin asymmetric combinations contribute to the latter, breaking the proportionality to the m^4 coefficient. The leading disorder contributions to the fluctuation corrections are given by,

$$\begin{aligned} F_{\text{fl}}^{\text{dis}} &= \tilde{g} \tilde{\sigma}_c^2 \sum_{\nu, \nu'} \int_{\mathbf{k}_1 \mathbf{k}_2 \mathbf{k}_3} h_{\nu, \nu'}(\mathbf{k}_1, \mathbf{k}_2) \left\{ (-1)^{\nu+\nu'} \right. \\ &\times \chi_{\nu, \nu'}(\mathbf{k}_1, \mathbf{k}_2) \chi_{-\nu, -\nu'}(\mathbf{k}_3, \mathbf{k}_3 - \mathbf{k}_1 + \mathbf{k}_2) \\ &\left. + 2K_{\nu, \nu'}(\mathbf{k}_1, \mathbf{k}_2) n_{-\nu'}(\mathbf{k}_3) \right\}, \end{aligned} \quad (7)$$

with $K_{\nu, \nu'}(\mathbf{k}_1, \mathbf{k}_2) = \tilde{T} \sum_{\omega_n} G_{\nu}(\mathbf{k}_1, \omega_n) G_{\nu'}^2(\mathbf{k}_2, \omega_n)$ [17].

To obtain the corrections to the phase boundaries we compute $F_{\text{mf}}^{\text{dis}}$ and $F_{\text{fl}}^{\text{dis}}$ numerically and extract the contributions to the m^2 , m^4 , and $q^2 m^2$ coefficients as functions of \tilde{T} and \tilde{g} . The resulting phase diagram is shown in Fig. 1. Disorder has two effects. On the one hand it destabilizes the homogeneous FM due to a positive contribution to the m^2 coefficient. On the other hand, it enhances the tendency towards spiral ordering because of a dominant, negative fluctuation correction to the




FIG. 2: (color online) Three-dimensional plots of (a) the spiral ordering wave vector, (b) correlation length, and (c) anisotropy across the helical spin glass phase.

$q^2 m^2$ term. While small disorder leads to a slight increase of the tricritical temperature \tilde{T}_c , this effect is reversed by higher-order contributions that cut-off the non-analyticity of the clean system, $F_{\text{fl}} \sim \tilde{g}^2 m^4 \ln[m^2 + (\tilde{T} + (k_F l)^{-1})^2]$ (l the mean free path) and lead to a new non-analytic term of opposite sign [6, 18]. Therefore, for sufficiently strong disorder, the tricritical point is pushed to lower temperatures and is eventually destroyed.

Disordered XY-Phenomenology in the Helical Spin Glass: Charge disorder does not destroy long range order (LRO) in the homogeneous FM. This is captured appropriately by our analysis of the replica averaged free energy. To see this, we may formally introduce independent order parameters \mathbf{m}_α in each replica. The resulting $\mathbf{m}_\alpha^2 \mathbf{m}_\beta^2$ and $(\nabla \mathbf{m}_\alpha)^2 (\nabla \mathbf{m}_\beta)^2$ terms would arise from a replica average of disorder that couples to \mathbf{m}^2 and $(\nabla \mathbf{m})^2$, respectively. Such rotationally symmetric random mass and stiffness disorder does not destroy LRO in the FM.

To understand the relevance of disorder in the fluctuation driven spiral region we must analyze how disorder couples to the Goldstone modes. Taking into account lattice effects, the free energy is no longer invariant under continuous rotations of the spiral ordering wave vector

q. Starting, for example, from a tight-binding dispersion on a cubic lattice and a chemical potential close to the bottom of the band, we obtain a small anisotropy term $F_{\text{anis}} = \lambda \int d^3\mathbf{r} [(\partial_x m_x)^2 + (\partial_y m_y)^2 + (\partial_z m_z)^2]$ ($\lambda > 0$), which is minimized for \mathbf{q} along one of the crystal axes. In the following we will assume $\mathbf{q} = q\mathbf{n}_z$ as before. Note that the anisotropy term fixes the direction of \mathbf{q} without changing the saddle-point equation $q^2 = -f_2(m^2)/f_4(m^2)$ for the modulus of \mathbf{q} . The Goldstone modes correspond to smooth deformations $\phi(\mathbf{r})$ of the phase of the spiral order parameter, $\mathbf{m}(\mathbf{r}) = m[\mathbf{n}_x \cos(qz + \phi) + \mathbf{n}_y \sin(qz + \phi)]$, captured by the classical anisotropic 3d-XY action

$$S_\phi = \int d^3\mathbf{r} \left\{ 2f_4(m^2)q^2(\partial_z \phi)^2 + \frac{1}{2}\lambda m^2 \sum_{i=x,y} (\partial_i \phi)^2 \right\}. \quad (8)$$

Disorder in the spin stiffness induces local random changes of q . While this obviously leads to disorder in the stiffness of the Goldstone modes it also generates a random anisotropy for the phase angle of the spiral,

$$S_\phi^{\text{dis}} = \frac{1}{2}\lambda m^2 \int d^3\mathbf{r} g(\mathbf{r}) \cos[2\phi + \alpha(\mathbf{r})], \quad (9)$$

with $\alpha(\mathbf{r}) \sim \delta q(\mathbf{r})z$ a random phase and $g(\mathbf{r}) = [(\partial_y \alpha)^2 - (\partial_x \alpha)^2]/4$. No such random anisotropy is induced in the homogenous FM phase, leading to a very different response of the spiral phase to disorder. In particular, it is established that arbitrarily small disorder of random field or higher-order random anisotropy type destroys LRO in dimensions $d < 4$ [19–21]. It is very likely that the random anisotropy 3d-XY model - and so the spiral phase - supports only short-ranged magnetic correlations [22].

This phase of short-ranged magnetic order is markedly anisotropic. In the following, we use the result by Dotenko and Feigelman [23] who argued that on scales much larger than the size of a vortex-line core, the correlation length of the isotropic 3d-XY model with random anisotropy disorder, $D \cos(n\phi + \alpha(\mathbf{r}))$, and spin stiffness ρ_s is given by $\xi = \frac{n^2}{8\pi} \frac{\rho_s^2}{D^2 K_n^2}$, where $K_n(T) = \langle \cos(n\phi) \rangle_0$ is computed with the wave-like excitation in the absence of disorder. Specifying to $n = 2$ and taking into account a rescaling of the z -coordinate, we obtain

$$\xi_{xy} = \frac{1}{2\pi\sigma_g^2} \left(\frac{\lambda m^2}{4f_4(m^2)q^2} \right)^2 e^{\frac{\tilde{T}}{\pi\sqrt{4\lambda f_4(m^2)m^2 q^2}}} \quad (10)$$

for the correlation length along the directions perpendicular to \mathbf{q} where σ_g^2 is the variance of $g(\mathbf{r})$. The anisotropy is given by $\xi_z/\xi_{xy} = \sqrt{4f_4(m^2)q^2/(\lambda m^2)}$. Since from minimization of the free energy m and q are obtained as a function of temperature \tilde{T} and electron repulsion \tilde{g} we can evaluate $\xi = \xi(\tilde{T}, \tilde{g})$ over the spiral region. In Fig. 2 the evolution of q , $1/\xi_{xy}$, and ξ_z/ξ_{xy} is shown.

At the transition to the homogeneous FM, the magnetization remains finite while $q \rightarrow 0$, leading to a divergence of ξ . Note that in the presence of a magnetic easy axis anisotropy, this transition becomes weakly first order [24]. The correlation length changes significantly over the spiral region and is strongly anisotropic with $\xi_z > \xi_{xy}$, except very close to the transition to the FM. In the clean limit, $\sigma_g^2 \rightarrow 0$, LRO is recovered in the spiral phase.

Discussions and Conclusions – By combining a replica disorder average with the fermionic quantum order-by-disorder approach [10] we have demonstrated that weak charge disorder destabilizes the FM phase but enhances the susceptibility towards incommensurate spiral ordering below the tricritical point. While disorder does not destroy LRO in the FM phase, the correlation length in the spiral region is finite even for infinitesimal disorder. This is a consequence of the interplay of the (cubic) crystal anisotropy and disorder in the spin-stiffness which generates a relevant random anisotropy term in the effective 3d-XY model for the Goldstone modes.

A helical glass with short-range order is consistent with the peculiar incommensurate state recently found close to the avoided FM quantum critical point of CeFePO [13]. The μ SR results indicate that the sample-averaged muon asymmetry function is dominated by strongly inhomogeneous spin fluctuations and shows the characteristic time-field scaling expected from glassy spin dynamics. The anisotropy of magnetic fluctuations and the lack of evidence for FM cluster formation in field cooled magnetization measurements rule out conventional spin-glass behavior [13]. Both observations find a natural explanation in terms of short range spiral order.

The presence of oriented vortex-line defects in the helical glass should lead to a non-Fermi-liquid $T^{3/2}$ temperature dependence of the electrical resistivity, as observed in the partially ordered phase of the helimagnet MnSi [25, 26]. From a symmetry point of view, helical magnetic phases are analogous to the ordered phases of cholesteric liquid crystals [14] and the partial magnetic order in MnSi is reminiscent of blue phases of cholesterics [26], where the line defects correspond to columnar skyrmion spin textures [15]. The vortex lines that we discuss here are rather different from skyrmion line defects. The latter correspond to vortices in the orientation of the spiral wave-vector, whereas the vortices predicted here are in the pitch of the spiral order for a fixed direction of wave vector. Nevertheless, some properties of line defects are quite generic. In Ref. [27], it was shown that columnar fluctuations of any type of line defects lead to a $T^{3/2}$ contribution to the electrical resistivity. This mechanism should equally apply to the extended phase defects that we discuss here.

The authors benefited from stimulating discussions with M. Brando A. V. Chubukov, and G. J. Conduit. This work was supported by EPSRC under grant code EP/H049584/1. SJT acknowledges financial support

from the Scottish CM-DTC.

-
- [1] C. Pfleiderer, S. R. Julian, and G. G. Lonzarich, *Nature* **414**, 427 (2001).
- [2] Y. J. Uemura, T. Goko, I. M. Gat-Malureanu, J. P. Carlo, P. L. Russo, A. T. Savici, A. Aczel, G. J. MacDougall, J. A. Rodriguez, G. M. Luke, et al., *Nat. Phys.* **3**, 29 (2007).
- [3] M. Otero-Leal, F. Rivadulla, M. Garca-Hernandez, A. Pineiro, V. Pardo, D. Baldomir, and J. Rivas, *Phys. Rev. B* **78**, 180415(R) (2008).
- [4] V. Taufour, D. Aoki, G. Knebel, and J. Flouquet, *Phys. Rev. Lett.* **105**, 217201 (2010).
- [5] E. A. Yelland, J. M. Barraclough, W. Wang, K. V. Kamenev, and A. D. Huxley, *Nat. Phys.* **7**, 890 (2011).
- [6] D. Belitz, T. R. Kirkpatrick, and T. Vojta, *Phys. Rev. Lett.* **82**, 4707 (1999).
- [7] A. V. Chubukov, C. Pépin, and J. Rech, *Phys. Rev. Lett.* **92**, 147003 (2004).
- [8] G. J. Conduit, A. G. Green, and B. D. Simons, *Phys. Rev. Lett.* **103**, 207201 (2009).
- [9] F. Krüger, U. Karahasanovic, and A. G. Green, *Phys. Rev. Lett.* **108**, 067003 (2012).
- [10] U. Karahasanovic, F. Krüger, and A. G. Green, *Phys. Rev. B* **85**, 165111 (2012).
- [11] T. Vojta, *Phys. Rev. Lett.* **90**, 107202 (2003).
- [12] J. A. Hoyos and T. Vojta, *Phys. Rev. Lett.* **100**, 240601 (2008).
- [13] S. Lausberg, J. Spehling, A. Steppke, A. Jesche, H. Luetkens, A. Amato, C. Baines, C. Krellner, M. Brando, C. Geibel, et al., *Phys. Rev. Lett.* **109**, 216402 (2012).
- [14] D. Belitz, T. R. Kirkpatrick, and A. Rosch, *Phys. Rev. B* **73**, 054431 (2006).
- [15] U. K. Rössler, A. N. Bogdanov, and C. Pfleiderer, *Nature* **442**, 797 (2006).
- [16] For the contact interaction we obtain $\lambda = 16\sqrt{2}/[3(2\pi)^6]$ and a tricritical point at $\tilde{T}_c \approx 0.3$ (see Fig. 1), consistent with earlier results [6]. With increasing range of the interaction, λ decreases considerably, leading to an exponential suppression of the tricritical temperature [6, 28]. κ is a constant arising from re-summation of the leading divergencies [29] and controls where the first-order transition terminates on the $\tilde{T} = 0$ axis. In this letter we use the value $\kappa = 0.001$.
- [17] Note that the matrix elements $K_{\nu,\nu'}$ can be written as combinations of derivatives of $\chi_{\nu,\nu'}$ with respect to the chemical potential and the magnetization.
- [18] T. R. Kirkpatrick and D. Belitz, *Phys. Rev. B* **85**, 134451 (2012).
- [19] Y. Imry and S. Ma, *Phys. Rev. Lett.* **35**, 1399 (1975).
- [20] R. A. Pelcovits, E. Pytte, and J. Rudnick, *Phys. Rev. Lett.* **40**, 476 (1979).
- [21] M. Aizenman and J. Wehr, *Phys. Rev. Lett.* **62**, 2503 (1989).
- [22] The more exotic possibility of algebraic quasi LRO seems unlikely. Near $d = 4$ an infinite number of operators become relevant, requiring a functional RG (FRG) treatment [30]. For sufficiently weak random anisotropy disorder as present in the spiral region, it has been demonstrated within 2-loop FRG that $O(N)$ magnets with $N < N_c = 9.44$ exhibit algebraic quasi long range order for $d_{lc} < d < 4$ with a lower critical dimension $d_{lc} \approx 4 - 0.00158(N - N_c)^2$ [31] which gives $d_{lc}(N = 2) \approx 3.91$ in the XY case. While such extrapolation should be taken with caution given the high value of N_c [31], very similar lower critical dimensions only slightly below $d = 4$ are found for random field disorder within non-perturbative FRG [32]. It is therefore very likely that the disordered phase in the random anisotropy $3d$ -XY model is conventional with short-range magnetic correlations.
- [23] V. S. Dotsenko and M. V. Feigelman, *J. Phys. C* **16**, L803 (1983).
- [24] G. Abdul-Jabbar, F. Krüger, D. Sokolov, C. Stock, F. Demmel, D. Wermeille, F. Levy, B. Grenier, A. G. Green, and A. Huxley, in preparation.
- [25] N. Doiron-Leyraud, I. R. Walker, L. Taillefer, M. J. Steiner, S. R. Julian, and G. G. Lonzarich, *Nature* **425**, 595 (2001).
- [26] C. Pfleiderer, D. Reznik, L. Pintschovius, H. v. Löhneysen, M. Garst, and A. Rosch, *Nature* **427**, 227 (2004).
- [27] T. R. Kirkpatrick and D. Belitz, *Phys. Rev. Lett.* **104**, 256404 (2010).
- [28] C. W. von Keyserlingk and G. J. Conduit, arXiv:1301.6036.
- [29] C. Pedder, F. Krüger, and A. G. Green, in preparation.
- [30] D. S. Fisher, *Phys. Rev. B* **31**, 7233 (1985).
- [31] P. Le Doussal and K. J. Wiese, *Phys. Rev. Lett.* **96**, 197202 (2006).
- [32] M. Tissier and G. Tarjus, *Phys. Rev. Lett.* **96**, 087202 (2006).